Uncommon 50-58 mm (2.0-2.3 in.) Medium

MALE: Head: prominent median spike on occiput. Face: pale yellow-green. Eyes: pale blue-gray/aqua-gray. Thorax: yellowish-green w/ middorsal black stripe w/ gray-green patches to either side. Abdomen: black w/ narrow light yellow pattern on S1-6 and gray ventrally; S7-9 black and swollen; S10 light bordered w/ black; lateral thick black line or thin black line w/ gray-green lines between and below. Abdomen: middorsally yellow. Legs: short w/ short black spines. Wings: clear. Stigmas: narrow, light brown. Appendages: light yellow, wide open and hooked.

FEMALE: Similar to M. (Genus is only Clubtail w/ an ovipositor.)

Habitat: Ponds (usually) or edges of slow-moving Coastal Plain rivers w/ muddy bottom and little submerged vegetation. Commonly perches on lily pads or muddy banks of slow steams or ponds. Tolerates degraded habitats.

Reproduction: Patroling: low over water. Female oviposits alone, tapping abdomen in water near shore. Exuviae at water edge or floating on the surface (high enough to free head/thorax).

Notes: Was Gomphus villosipes.

Locations: Centennial - VP, Cent HS, Elkhorn - L/FB, Font Hill, Gateway - RF/SM, Henryton, Homespun, Jackson Pond, Kittamaqundi (north end; RO), Manor Woods, Marriottsville, Meadowbrook, Milk Plant, MPEA P3/P4, Old Frederick, Schooley Mill, Sewell's Orchard, Sunnyside, Trinity, VFW Pond. Western Regional, Woodstock.

Phenograms (pages 2 – 6).

Locations (in Howard County)

Identification features and odonate anatomy